

Chemistry of Heterocyclic Compounds 1968 vol.2 N5, pages 493-495

---

## Reaction of cycloheptanone with concentrated sulfuric acid

Feller K., Svetozarskii S., Samitov Y., Zil'berman E., Razuvaev G.

*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

### Abstract

Treatment of cycloheptanone or 2-cycloheptylidencycloheptanone with concentrated sulfuric acid, leads to self-condensation accompanied by intramolecular rearrangement and cyclization to give substituted tetrahydrofurans, viz. 2-methyl-2, 3-cyclohexano-4, 5-cyclohepteno-( $\Delta$  5)-2, 3, 4, 5-tetrahydrofuran and 2-methyl-2, 3-cyclohexano-5-hydroxy-4, 5-cycloheptano-2, 3, 4, 5-tetrahydrofuran. Self-condensation of ketones with 6-membered rings under the action of concentrated sulfuric acid, proceeding via carbonium ions, and accompanied by contraction to a 5-membered ring, is also characteristic of cycloheptanone, the 7-membered ring of which isomerizes to a 6-membered one. © 1968 The Faraday Press, Inc.

<http://dx.doi.org/10.1007/BF00477501>

---